

# Retinex Theory

*Draft for Encyclopedia of Color Science and Technology, 2nd Edition, Springer*

**John McCann**

[mccanns@tiac.net](mailto:mccanns@tiac.net)

McCann Imaging, 30 Spy Pond Parkway, Arlington, MA 02474, USA

## Definition

Retinex is the theory of human color vision proposed by Edwin Land to account for color sensations in real scenes. Color constancy experiments showed that color does not correlate with receptor responses. In real scenes, the content of the entire image controls appearances. A triplet of L, M, S cone responses can appear any color. Land coined the word “Retinex” (the contraction of retina and cortex) to identify the spatial image processing responsible for color constancy. Further, he showed that color sensations are predicted by three apparent lightnesses observed in long-, middle-, and short-wave illumination. Retinex is also used as the name of computer algorithms that mimic vision’s spatial interactions to calculate the apparent lightnesses observed in complex scenes.

## Overview


Edwin H. Land, the inventor of hundreds of photographic film patents, was struck by experiments showing that color sensations in real complex images depend on scene content. Film responds to the light falling on each tiny local region. Land realized that vision’s mechanisms were very different from film. His early experiments studied the colors observed in red and white projections [ 1]. He realized color appearance required both the cone responses to a local region and the neural spatial processing of the rest of the scene. He proposed the Retinex Theory.

Land coined the word Retinex to describe three independent spatial channels. In 1964 he wrote: “We would propose that all of the receptors with maximum sensitivity to the long-waves in the spectrum, for example, operate as a unit to form a complete record of long-wave stimuli from objects being observed. For convenience of reference, let us call this suggested retinal-cerebral system a ‘retinex.’)” [ 2– 5] It is the word that describes the mechanisms that performs the comparison of scene information to create the array of sensations of L, M, S lightness in each of human vision’s three channels.

## Cone Quanta Catch

Visible light falls on objects that reflect some of it to the eye. Color vision depends on the spectrum of the illumination falling on an object and the spectrum of its reflectance. The product of these spectra describes the light coming to the eye. There are three types of cones in normal observers that are called L for long-wave-, M for middle-wave-, and S for short-wave-sensitive cones. The receptors’ spectral sensitivities multiplied by the light falling on the retina determines the L, M, S cone responses, namely, the “quanta catch” of the cones. The cones convert the quanta catch to nerve signals that pass through many spatial comparisons in the visual system. The cone quanta catch is the important transition from the physics of light to the physiology of vision. However, it is just the first step in the process.

Figure 1 illustrates a laboratory experiment that generates equal L, M, S quanta catches from different reflectance papers.



### Retinex Theory, Fig. 1

Equal quanta catches from red and green papers. In a darkroom with a black surround, identical quanta catches appear the same - neither red, nor green color appearances. When viewed alone, in a no-light background, the appearance of a spot of light correlates with cone quanta catch.

At the top of Fig. 1, light passes through filters that determine the spectrum of each illumination falling on circular pieces of paper. The light coming to the eye from the papers is modified again by the reflectance spectra of the papers. Further, in this experiment there is no light coming from the black surrounds.

On the left, there is a tungsten light source at the top. There is a filter that absorbs more middle-wave than long- and short-wave visible light (magenta arrows). The green circular paper reflects more middle-wave than long- and short-wave light. The light coming to the eye is the product of these spectra. The integrals of that light using the three cone spectral sensitivities determine the L, M, S quanta catch values.


On the right, there is the same tungsten light source at the top. There is a different filter that absorbs more long-wave than middle- and short-wave light (cyan arrows). The right-red paper reflects more long-wave than middle- and short-wave light. In this experiment, the spectra of the two illuminants and the two reflectances were adjusted to generate the same triplet of LMS cone quanta catches. Under these conditions, the left-green paper and the right-red papers are identical retinal stimuli. They appear equal to each other, but that color is neither red nor green.

## Mondrian Experiments

Land's color Mondrian experiment is similar, with the exception that he used a complex array of papers to simulate real-world scenes. The important difference is that in complex scenes, a particular quanta catch can appear any color: red, green, blue, yellow, white, or black.

Figure 2 shows the double color Mondrian experiment [ 6]. It used two identical Mondrians made of color papers, and two identical sets of illuminators. Each set had three different, non-overlapping spectral illuminants (long-, middle-, and short-wave visible light). Each projector provided spatially uniform illumination for one spectral band, for one of the Mondrians. The amount of illumination from each projector was independently adjusted.

In this experiment observers reported the colors of papers in the Mondrians. In this illustration, we will look at the circular red and green papers. Land adjusted the illuminant mixtures of light from the two sets of three projectors in order to control the light coming from the green and red papers. The amounts of L, M, S light came from the green circle on the left as from the red circle on the right. First, he turned on just the long-wave lights. He adjusted the amounts of spatially uniform illumination falling on each Mondrian separately. He set the illumination so that the left-green circle and right-red circle had equal light meter readings. Then, he did the same for middle- and short-wave light. The left-green and right-red circles had equal quanta catches by the L, M, S cones.



**Retinex Theory, Fig. 2**


Land's double Mondrian experiment. Two identical sets of matte colored papers, with separate L, M, S illuminating projectors with voltage transformers for control of the amount of spatially uniform light falling on each Mondrian. Telephotometer readings were projected above the Mondrians. Land separately measured the L, M, S radiances from a green circle in the left Mondrian. Then, he adjusted the L, M, S radiances from a red circle on the right Mondrian to be the same. Observers reported different red and green colors produced by identical light stimuli

In this complex scene, observers reported that equal quanta catches circles appeared green on the left and red on the right. Observers reported color constancy, namely, that the red paper looked red and the green paper looked green despite the identical cone quanta catches.

Land repeated this experiment with all the Mondrian papers. A constant L, M, S quanta catch could generate any color sensation. The presence of the complex scene introduced more information to the visual system. The red and green papers appeared equal in Fig. 1; neither red, nor green. The red paper looked red and the green paper looked green in Fig. 2. The scene's spatial content stimulated vision's spatial image processing mechanisms to generate color constancy. The post-receptor visual processing plays a dominant role in color appearance in real scenes. Land's word "Retinex" gave this spatial process a name. As well, he proposed a theoretical mechanism.

## Retinex Mechanism

Figure 3 illustrates the pair of Mondrians in only long-wave light with more illumination on the left Mondrian, than on the right. The appearances of the two Mondrians are nearly constant. This is a common observation: humans are insensitive to large changes in uniform illumination.




### Retinex Theory, Fig. 3

Pair of Mondrians in only long-wave light. Each rectangle in the left Mondrian has the identical reflectance as the corresponding rectangle in the right Mondrian. Only the circles have different reflectances. The left Mondrian has more illumination than the right. Observers report that the left Mondrian rectangles appear slightly lighter than the right rectangles. Each corresponding rectangle is nearly the same lightness in the left and right Mondrians.

As illustrated here, in long-wave illumination the left-green paper circle appeared dark when it generated the same L cone quanta catch as the lighter right-red circle. Vision's spatial image processing rendered the red and green papers with different apparent lightnesses in long-wave light. The lightnesses are stable with large changes in overall illumination.

Figure 4 illustrates the Mondrian viewed in middle-wave illumination. The left-green circle now appeared light and the right-red circle looked dark. In Fig. 4 the circles had the same M cone quanta catch. Vision's spatial image processing rendered the red and green papers with different, and opposite lightnesses in Fig. 3 and Fig. 4.



#### Retinex Theory, Fig. 4

Pair of Mondrians in only middle-wave light. Now, the left Mondrian has less illumination than the right. In this waveband the pattern of lightnesses differs from that in long-wave light. Observers report that the right Mondrian rectangles are slightly lighter than the left ones. Each corresponding rectangle is nearly the same lightness in the left and right Mondrians

The left-green circular paper has more long-wave and less middle-wave illumination. The right-red circular paper has less long-wave and more middle-wave illumination. When adjusted using the telephotometer, those illuminations make the red and green papers have identical radiances. Using the same procedure Land adjust the short-wave illuminants to make the left and right circles to have equal radiances. Having set the L, M, S illuminations so that both circles have equal radiances, Land turned on all six projectors. Although both circles caused identical cone quanta catches, observers reported that they had very different color appearances; red and green. (illustrated in Fig 2) Using this procedure Land showed that a particular L, M, S quanta catch can appear any color.

James Clerk Maxwell invented Colorimetry based on color matches. Also, he invented Color Photography based on three-color-separations with silver halide photographs. Both Colorimetry matches and silver halide photography are molecular quanta catch events. It is easy to think that human vision uses those same mechanisms. But, color constancy disproves that hypothesis. Edwin Land's Retinex invention adopted the cone quanta catch as vision's initial

receptor response, but realized that vision's color appearance separations are the result of neural spatial comparisons, not receptors' cone quanta catch. Spatial comparisons are the basis of color constancy in complex scenes. Spatial comparisons form vision's L, M, S color separations.

These constancy observations led Land to propose the Retinex theory. The triplet of apparent lightnesses, not cone quanta catches, determines the color appearance. Constant LMS lightnesses generate constant colors. That hypothesis led to a study of color appearances in L, M, S bands of light. Do all red colors have the same triplet of lightness appearances? Does a red color always look [light, dark, dark] in L, M, S light? Does a green always look [dark, light, dark]? Does color appearance always correlate with the triplet of L, M, S lightnesses?

The experiment is easy. Find narrowband red, green, and blue filters. Be sure that the filters exclude the other two-thirds of the spectrum. With a green filter you should just see greens with different lightnesses. You should not see a mixture of greens and yellows and blues hues. If you do, you need a filter with a narrower band of transmission.

Identify a group of red objects. Look at them sequentially through the L, M, S filters. Look at them in different ambient illuminations. Look at them at different times of the day. Look at them in sunlight and shadows. Red colors are always (light, dark, dark) in L, M, S light. The same dependence of the triplet of L, M, S lightnesses holds for all colors (Table 1). Lightness is the output of spatial image processing. It is the result of post-receptor spatial processing. That is why lightness does not correlate with cone quanta catch. However, color does correlate with three lightnesses in long-, middle-, and short-wave light.

**Retinex Theory, Table 1**


Correlation table of color appearances and the apparent lightnesses in L, M, S illumination

Color Appearance	Appearance in L- light	Appearance in M- light	Appearance in S- light
Red	light	dark	dark
Yellow	light	light	dark
Green	dark	light	dark
Cyan	dark	light	light
Blue	dark	dark	light
Magenta	light	dark	light
White	light	light	light
Black	dark	dark	dark

Retinex theory predicts that the triplet of L, M, S lightnesses determines color. Colors are constant with changes in illumination because the triplet of lightnesses is nearly constant.

Land's observation still stands: The triplet of lightnesses correlates with color. The observation is important because a variety of different phenomena can influence lightness, such as simultaneous contrast, the Cornsweet effect, and assimilation. Regardless of the cause of the lightness changes, when two identical physical objects look different, color appearances correlate with their L, M, S lightnesses ([ 7], Section E).

In the color assimilation display (Fig. 5), there are two sets of nine red squares that have the same reflectance and appear the same (top left). However, if these red squares are surrounded by yellow and blue stripes, they look different (top center): the left red squares fall on top of the yellow stripes, and the right ones on the blue stripes. The left squares appear a purple red, while the right ones appear a yellow orange. In other words, the left squares appear more blue and the right ones more yellow.



### Retinex Theory, Fig. 5

( *Top-left*) Red color squares on white and ( *top-center*) color assimilation experiments change the appearance of the red squares; ( *bottom*) L, M, S separation images. Color appearances of *red squares* correlate with L, M, S lightnesses

In Fig. 5 (bottom–left) the L separation the corresponding squares are lighter on the right of the separation; in the M separation these patches are lighter; in the S separation they are darker on the right. Land’s Retinex predicts that whenever L and M separations are lighter and S separation is darker, then that patch will appear more yellow (See right squares in Assimilation experiment (top-center)]. Whenever S separation is lighter and L and M separations are darker, then that patch will appear more blue. [See left squares in Assimilation experiment (top-center)]. Colors correlate with L, M, S lightnesses ([ 7], pp. 221–281).

## Retinex Image Processing

Land described that the fundamental challenge of color vision shifted to the ability to predict lightness; that is, the spatial interactions in each color channel found in post-receptor neural processes. In 1967 Land and McCann proposed a computational model for calculating lightness from the array of all scene radiances [ 6]. The model compared each pixel with every other pixel in an image. The goal was to calculate the sensation of image segments that equaled what observers saw. In the past 50 years, there have been many implementations and variations of this process. They are called Retinex algorithms. It is curious that Land reserved the use of the term “Retinex” to describe three independent lightness channels. Today’s usage of the word includes a much wider range of computer algorithms that build calculated appearances out of arrays of radiances.

To calculate lightnesses in complex scenes, one must:

- Capture radiances from the entire scene.
- Convert scene radiances to cone and rod quanta catches.
- Calculate lightness for each color channel using all pixels in the scene.
- Compare calculated lightnesses with observer matches.

The Land and McCann model [ 6, 9, 11. ( 7 pp. 293-337)] used:

- Edge ratios of cone quanta catches (basic building block)
- Gradient threshold (found to be unnecessary in later studies)
- Multiplication of edge ratios (makes long-distance interactions)
- Reset to maxima (scaled the output)
  - (introduced dependence on scene content, e.g., simultaneous contrast)
- Average of many spatial comparisons

The first computer implementation of the model in the early 1970’s used an array of 20 by 24 pixels. McCann, McKee, and Taylor showed that long-, middle-, and short-wave computed lightnesses predicted observer matches of color Mondrians in color constancy experiments [ 8].

Since the late 1960s, computer imaging has shown remarkable advances. Digital images

have replaced film in most of photography. Computer graphics has made image synthesis ubiquitous. Retinex image processing has grown with the advances in digital imaging [ 9– 11]. In the early 1980s Frankle and McCann introduced a multi-resolution algorithm that allowed efficient comparison of all pixels in a 512 by 512 pixel image [ 12]. Multiresolution software programs (Zoom Retinex) calculates all scene lightnesses with  $O(N)$  computational efficiency [ 9]. Sobol extended Frankle and McCann’s algorithm in the design of commercial digital cameras [ 13] introduced in 2003. Other applications used Retinex’s efficient spatial processing for color gamut-mapping [ 14].

Using different approaches, Jobson, and Kotera have studied the NASA Retinex. Rizzi and colleagues have developed the Milan Retinex ([ 7], pp. 324–328).

The important feature of real complex scenes is that the illumination is rarely uniform. Shadows and multiple reflections increase the dynamic range of light coming to our eyes, and to cameras. The application of Retinex algorithms to high dynamic range (HDR) scenes has become a major topic of research and engineering applications ([ 7], pp. 283–375). The limits of HDR scene capture and reproduction are controlled by optics, namely, optical veiling glare. Camera glare limits the range of light on the sensor, just as intraocular glare limits the range of light on the retina. The scene content controls the distribution of glare range of light in images. Vision’s post-receptor neural processes compensate for veiling glare ([ 7], pp. 89–171). That explains humans’ high dynamic range of appearances from low-dynamic-range retinal images. The spatial mechanisms modeled by Retinex algorithms play a major role in compensating for glare and generating our range of color and lightness sensations [ 15, 16].

Over the years many variations of spatial processing mimicking human vision have been called Retinex algorithms.

## Signatures of Color Constancy

Studies of Color constancy have documented its limits. Measurements of departures from perfect constancy provide a signature of the underlying post-quantum-catch neural mechanisms. Land’s double Mondrian experiments used spatially uniform illumination with different spectral compositions. In each spectral illuminant observers chose matching Munsell chips very close to the actual Munsell papers in the Mondrian areas; indicating near perfect constancy. Nevertheless, each spectral illuminant had different small systematic departures from perfect constancy across the entire Mondrian [ 8]. Similar small departures were observed in experiments with color adapting surrounds, again in uniform spatial illumination ([ 7] pp. 239-245). A pair of recent experiments measured color constancy using two identical 3-D Mondrians made of blocks of wood painted with one of 11 matte paints. The experiments used different illuminations. The first used a diffusing tent that minimized shadows - the Low-Dynamic-Range (LDR) scene. The second used spotlights that made sharp shadows - the High-Dynamic-Range (HDR) scene. Observers reported a marked loss of color constancy in 3-D Mondrians. LDR illumination caused a substantial loss of constancy in some local regions; HDR illumination caused a much larger loss in other regions [ 17]. The appearance of a surface in 3-D blocks no longer shows universal correlation with the paint’s surface reflectance when the illumination introduces quanta-catch edges. These and other experiments support Retinex’s spatial comparison mechanism [ 18]. Five different studies of the signatures of color appearance in real scenes demonstrate the important role of spatial comparisons in vision [19].

## Summary

Edwin Land coined the term Retinex in 1964 as the name of human color channels that build appearance up from spatial comparisons. Over the years others have redefined the term to include many different spatial algorithms that attempt to mimic human spatial color processing. Land’s Retinex was built to model color constancy. Over the years quantitative measurements of the limits of constancy have provided signatures of the underlying visual mechanism. These signatures supports neural spatial comparisons within color channels.



## References

- [ 1] McCann, J.J., Benton, J., McKee, S.: Red/white projections and rod/long-wave cone color: an annotated bibliography. *J. Electron. Imaging* 13, 8–14 (2004)
- [ 2] Land, E.H.: The retinex. *Am. Sci.* 52, 247–264 (1964)
- [ 3] Land, E.H., McCann, J.J.: Method and system for reproduction based on significant visual boundaries of original subject, U.S. Patent 3,553,360, June 5, 1971.
- [ 4] Land, E.H.: The retinex theory of colour vision. *Proc. R. Inst Gr. Brit.* 47, 23–58 (1974)
- [ 5] Land, E.H.: The retinex theory of color vision. *Sci. Am.* 237, 108–128 (1977)
- [ 6] Land, E.H., McCann, J.J.: Lightness and retinex theory. *J. Opt. Soc. Am.* 61, 1–11 (1971)
- [ 7] McCann, J., Rizzi, A.: *The Art and Science of HDR Imaging*, pp. 221–375. Wiley, Chichester (2012)
- [ 8] McCann, J.J., McKee, S., Taylor, T.: Quantitative studies in retinex theory, a comparison between theoretical predictions and observer responses to color Mondrian experiments. *Vision Res.* 16, 445–458 (1976)
- [ 9] McCann, J.J.: Lessons learned from Mondrians applied to real images and color gamuts. In: *Proceedings of IS&T/SID Color Imaging Conference*, vol. 7, pp. 1–8. Scottsdale (1999)
- [ 10] McCann, J.J.: Simultaneous Contrast and Color Constancy: Signatures of Human Image Processing, Chapter 6 in *Color Perception: Philosophical, Psychological, Artistic, and Computational Perspectives*, Vancouver Studies in Cognitive Studies: Davis, A., Davis, S. (eds.) pp. 87–101. Oxford University Press, Vancouver 2000
- [ 11] McCann, J.J.: Capturing a black cat in shade: past and present of retinex color appearance models. *J. Electron. Imaging* 13, 36–47 (2004)
- [ 12] Frankle, J., McCann, J.: Method and apparatus of lightness imaging. US Patent 4,384,336 (1983)
- [ 13] Sobol, R.: Improving the retinex algorithm for rendering wide dynamic range photographs. *J. Electron. Imaging* 13, 65–74 (2004)
- [ 14] McCann, J.J.: A spatial color-gamut calculation to optimize color appearance. In: MacDonald, L., Luo, R. (eds.) *Colour Image Science: Exploiting Digital Media*, pp. 213–233. Wiley, Chichester (2002)
- [ 15] McCann, J.J., V. Vonikakis, V., Rizzi, A.: *HDR Scene Capture and Appearance*. SPIE Spotlight Tutorial, Bellingham (2018)  
<[https://spie.org/Publications/Book/2315540?&origin\\_id=x109925&SSO=1](https://spie.org/Publications/Book/2315540?&origin_id=x109925&SSO=1)>
- [ 16] McCann, J.J., V. Vonikakis, V.: Calculating Retinal Contrast from Scene Content: A Program. *Front. Psychol.*, 17 Jan (2018)  
<<https://www.frontiersin.org/articles/10.3389/fpsyg.2017.02079/full>>
- [ 17] McCann, J.J., Parraman, C., Rizzi, A.: Reflectance, illumination, and appearance in color constancy, *Frontiers in Psychology*, 24, January 2014, 00005  
<<http://dx.doi.org/10.3389/fpsyg.2014.00005>>

[ 18] McCann, J.J.: Retinex at 50: color theory and spatial algorithms, a review. J.Electron. Imaging **26**(3), 031204 (2017), <<https://doi:10.1117/1.JEI.26.3.031204>>

[ 19]. McCann, J.J.: Limits of Color Constancy: Comparison of signatures of chromatic adaptation and spatial comparisons. In Proc. Electronic Imaging: Color Imaging XXIV, IS&T, San Francisco, (2019) <<https://doi.org/10.2352/ISSN.2470-1173.2019.14.COLOR-085>>